

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/495, 9/08, 47/40, 47/10	A1	(11) International Publication Number: WO 95/08993 (43) International Publication Date: 6 April 1995 (06.04.95)
(21) International Application Number: PCT/EP94/03169 (22) International Filing Date: 22 September 1994 (22.09.94) (30) Priority Data: 129,504 30 September 1993 (30.09.93) US (60) Parent Application or Grant (63) Related by Continuation US 129,504 (CIP) Filed on 30 September 1993 (30.09.93) (71) Applicant (for all designated States except US): JANSSEN PHARMACEUTICA N.V. [BE/BE]; Turnhoutseweg 30, B- 2340 Beerse (BE). (72) Inventors; and (75) Inventors/Applicants (for US only): FRANÇOIS, Marc, Karel, Jozef [BE/BE]; Valère Brockaertstraat 53, B-1090 Brussel (BE). DRIES, Willy, Maria, Albert, Carlo [BE/BE]; Molen- zijde 17, B-2330 Merksplas (BE). (74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica N.V., Patent Dept., Turnhoutseweg 30, B-2340 Beerse (BE).		(81) Designated States: AM, AU, BB, BG, BR, BY, CA, CN, CZ, FI, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LV, MD, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: ORAL FORMULATIONS OF AN ANTIFUNGAL (57) Abstract The present invention concerns a formulation for oral administration comprising an antifungal, a sufficient amount of a cyclodextrin or a derivative thereof, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent. Addition of one or more pharmaceutically acceptable sweeteners and one or more pharmaceutically acceptable flavours thereto yields palatable oral formulations. A process of preparing such formulations and pharmaceutical dosage forms comprising said formulations.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

ORAL FORMULATIONS OF AN ANTIFUNGAL

5

The present invention is concerned with novel compositions of antifungal agents which have low solubility in aqueous media, a process for preparing said compositions and pharmaceutical dosage forms for oral administration comprising said novel
10 compositions.

The development of efficacious pharmaceutical compositions of azole antifungals such as for example, itraconazole and saperconazole, is hampered considerably by the fact that said antifungals are only very sparingly soluble in water. The solubility and
15 bioavailability of said compounds can be increased by complexation with cyclodextrins or derivatives thereof as described in WO 85/02767 and US-4,764,604. Alternatively, strongly acidic formulations ($\text{pH} \leq 1.5$) of itraconazole and saperconazole can be formed in which the active ingredients are partially dissolved. Obviously such strongly acidic formulations are useless for oral administration. Aqueous formulations comprising a co-
20 solvent such as PEG 400 completely dissolve itraconazole at pH 2.3 -2.5. However, these acidic formulations have problems with regard to ease-of-preparation, acceptability, palatability and especially bioavailability : upon administration said formulations can precipitate irreversibly, e.g. in the stomach. Acidic formulations comprising cyclodextrin or a derivative thereof might appear an obvious alternative, but
25 the mere combinations prove to suffer from a number of similar problems, in particular difficulty-of-preparation, lack of stability (shelf life) and palatability, and unreliable absorption. In short, there still exists an important demand for easily prepared formulations of antifungal agents with good bioavailability and acceptable organoleptic properties for oral administration.

30

The present invention relates to formulations for oral administration which comprise an antifungal, e.g. itraconazole or saperconazole, as active ingredient, a sufficient amount of a cyclodextrin or a derivative thereof as a solubilizer, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent that greatly simplifies the preparation of
35 the composition. Preferred formulations are rendered more palatable by adding one or more pharmaceutically acceptable sweeteners, and one or more pharmaceutically acceptable flavours.

-2-

A low-dosage formulation according to the present invention is suitable for treating patients suffering from fungal infections, particularly for treating AIDS patients with oral candidiasis infections. The need for reliable formulations of itraconazole (and
5 saperconazole) in this indication is especially high because of resistance to fluconazole developing in *Candida* strains. Generally, 400 mg/day represents the minimum dose required to obtain meaningful plasma levels. Suitable oral formulations typically comprise from about 0.5% to about 1.5% (w/v), preferably about 1% (w/v) of the active
10 ingredient.

A high-dosage formulation according to the present invention is suitable for treating patients suffering from systemic fungal infections. Suitable oral formulations for combating systemic fungal infections typically comprise from about 3% to about 5%, preferably about 4% (w/v) of the active ingredient.

15 The formulations of the present invention are also suitable for the treatment of fungal infections in non-human animals, in particular for the treatment of dermatophytoses.

Itraconazole or (+)-cis-4-[4-[4-[4-[[2-(2,4-dichlorophenyl)-2-(1H-1,2,4-triazol-1-yl)-methyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]phenyl]-2,4-dihydro-2-
20 (1-methylpropyl)-3H-1,2,4-triazol-3-one, is a broadspectrum antifungal compound developed for oral, parenteral and topical use and is disclosed in US-4,267,179. Its difluoro analog, saperconazole or (\pm)-cis-4-[4-[4-[4-[[2-(2,4-difluorophenyl)-2-(1H-1,2,4-triazol-1-yl)methyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-1-piperazinyl]-phenyl]-
25 2,4-dihydro-2-(1-methoxypropyl)-3H-1,2,4-triazol-3-one, has improved activity against *Aspergillus* spp. and is disclosed in US-4,916,134.

Appropriate cyclodextrin derivatives are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units
30 of the cyclodextrin are substituted with C₁₋₆alkyl, particularly methyl, ethyl or isopropyl; hydroxyC₁₋₆alkyl, particularly hydroxyethyl, hydroxypropyl or hydroxybutyl; carboxyC₁₋₆alkyl, particularly carboxymethyl or carboxyethyl; C₁₋₆alkyl-carbonyl, particularly acetyl; C₁₋₆alkyloxycarbonylC₁₋₆alkyl or carboxyC₁₋₆alkyloxyC₁₋₆alkyl, particularly carboxymethoxypropyl or carboxyethoxypropyl; C₁₋₆alkyl-carbonyloxyC₁₋₆alkyl, particularly 2-acetyloxypropyl. Especially noteworthy as
35 complexants and/or solubilizers are β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- γ -CD, 2-hydroxypropyl- γ -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD.

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxypropyl and hydroxyethyl.

5 The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. In the cyclodextrin derivatives for use in the compositions according to the present invention the M.S. is in the range of 0.125 to 10, in particular of 0.3 to 3, or from 0.3 to 1.5. Preferably the M.S. ranges
10 from about 0.3 to about 0.8, in particular from about 0.35 to about 0.5 and most particularly is about 0.4. M.S. values determined by NMR or IR preferably range from 0.3 to 1, in particular from 0.55 to 0.75.

The average substitution degree (D.S.) refers to the average number of substituted
15 hydroxyls per anhydroglucose unit. In the cyclodextrin derivatives for use in the compositions according to the present invention the D.S. is in the range of 0.125 to 3, in particular of 0.2 to 2 or from 0.2 to 1.5. Preferably the D.S. ranges from about 0.2 to about 0.7, in particular from about 0.35 to about 0.5 and most particularly is about 0.4. D.S. values determined by NMR or IR preferably range from 0.3 to 1, in particular from
20 0.55 to 0.75.

More particular β - and γ -cyclodextrin hydroxyalkyl derivatives for use in the compositions according to the present invention are partially substituted cyclodextrin derivatives wherein the average degree of alkylation at hydroxyl groups of different
25 positions of the anhydroglucose units is about 0% to 20% for the 3 position, 2% to 70% for the 2 position and about 5% to 90% for the 6 position. Preferably the amount of unsubstituted β - or γ -cyclodextrin is less than 5% of the total cyclodextrin content and in particular is less than 1.5%. Another particularly interesting cyclodextrin derivative is randomly methylated β -cyclodextrin.

30 Most preferred cyclodextrin derivatives for use in the present invention are those partially substituted β -cyclodextrin ethers or mixed ethers having hydroxypropyl, hydroxyethyl and in particular 2-hydroxypropyl and/or 2-(1-hydroxypropyl) substituents.

35 The most preferred cyclodextrin derivative for use in the compositions of the present invention is hydroxypropyl- β -cyclodextrin having a M.S. in the range of from 0.35 to 0.50 and containing less than 1.5% unsubstituted β -cyclodextrin. M.S. values determined by NMR or IR preferably range from 0.55 to 0.75.

-4-

Substituted cyclodextrins can be prepared according to procedures described in US-3,459,731, EP-A-0,149,197, EP-A-0,197,571, US-4,535,152, WO-90/12035 and GB-2,189,245. Other references describing cyclodextrins for use in the compositions according to the present invention, and which provide a guide for the preparation, purification and analysis of cyclodextrins include the following : "Cyclodextrin Technology" by József Szejtli, Kluwer Academic Publishers (1988) in the chapter Cyclodextrins in Pharmaceuticals; "Cyclodextrin Chemistry" by M.L. Bender et al., Springer-Verlag, Berlin (1978); "Advances in Carbohydrate Chemistry", Vol. 12 Ed. by M.L. Wolfrom, Academic Press, New York (1975) in the chapter The Schardinger Dextrins by Dexter French at p. 189-260; "Cyclodextrins and their Inclusions Complexes" by J. Szejtli, Akademiai Kiado, Budapest, Hungary (1982); I. Tabushi in Acc. Chem. Research, 1982, 15, p. 66-72; W. Sanger, Angewandte Chemie, 92, p. 343-361 (1981); A. P. Croft and R. A. Bartsch in Tetrahedron, 39, p. 1417-1474 (1983); Irie et al. Pharmaceutical Research, 5, p. 713-716, (1988); Pitha et al. Int. J. Pharm. 29, 73, (1986); DE 3,118,218; DE-3,317,064; EP-A-94,157; US-4,659,696; and US-4,383,992. The low-dosage oral formulations according to the present invention typically comprise from about 20% to about 60% (w/v), preferably about 40% (w/v) of the cyclodextrin. The high-dosage formulations typically comprise from about 50% to about 80% (w/v), preferably about 60% (w/v) of the cyclodextrin derivative.

In order to increase the rate of dissolution of the poorly soluble antifungal during the manufacturing process, an alcoholic co-solvent is employed in the formulations according to the present invention. For this purpose, preference is given to those alcoholic co-solvents that have good dissolving power for itraconazole and/or saperconazole, in particular ethanol, propylene glycol and glycerol, especially propylene glycol. Without the alcoholic co-solvent, the dissolution of itraconazole or saperconazole in an aqueous acidic cyclodextrin medium is very slow, requiring a viscous suspension to be stirred for a prohibitively long time until complete dissolution is obtained. Addition of the alcoholic co-solvent, in the range of about 1% (v/v) to about 20% (v/v), preferably about 10% (v/v), increases the dissolution rate of the antifungal agent in an aqueous acidic cyclodextrin medium by a factor of at least 5 (when used at 10% (v/v)) and thus considerably shortens and simplifies the production process.

As a bulk liquid carrier there is used an acidic aqueous medium. Preferably the acidity of said carrier derives from a strong, pharmaceutically acceptable acid such as hydrochloric acid. The bioavailability of the antifungal agent and the organoleptic properties of the oral formulations are affected contrariwise by the acidity. An optimum effect can be

-5-

obtained at $\text{pH } 2.0 \pm 0.1$: that is, at this pH value, a sufficiently stable and bioavailable antifungal formulation is obtainable, the organoleptic properties of which can be rendered acceptable.

- 5 Not surprisingly, the ingredients thus far described yield a fairly strong-tasting potion when mixed with one another. Besides the acid taste due to the low pH, a bitter taste originating from the active ingredient, and possibly from the co-solvent (e.g. in the case of propylene glycol), is also present. Optimum taste masking can be obtained by the use of two types of adjuvants, namely pharmaceutically acceptable sweeteners and flavours.
- 10 Sweeteners are the more important additives in the low-dosage formulations, whereas the flavours are more important in the high-dosage formulations.

The pharmaceutically acceptable sweeteners comprise preferably at least one intense sweetener such as saccharin, sodium or calcium saccharin, aspartame, acesulfame potassium, sodium cyclamate, alitame, a dihydrochalcone sweetener, monellin, 15 stevioside or sucralose (4,1',6'-trichloro-4,1',6'-trideoxygalactosucrose), preferably saccharin, sodium or calcium saccharin, and optionally a bulk sweetener such as sorbitol, mannitol, fructose, sucrose, maltose, isomalt, glucose, hydrogenated glucose syrup, xylitol, caramel or honey.

20 The intense sweetener is conveniently employed in low concentrations. For example, in the case of sodium saccharin, the concentration may range from 0.04% to 0.1% (w/v) based on the total volume of the final formulation, and preferably is about 0.06% in the low-dosage formulations and about 0.08% in the high-dosage ones. The bulk sweetener 25 can effectively be used in larger quantities ranging from about 10% to about 35%, preferably from about 10% to 15% (w/v). In the high-dosage formulations the cyclodextrin derivative behaves as a bulk sweetener and none of the aforementioned bulk sweeteners needs to be added.

30 The pharmaceutically acceptable flavours which can mask the bitter tasting ingredients in the low-dosage formulations are preferably fruit flavours such as cherry, raspberry, black currant or strawberry flavour. A combination of two cherry flavours was found to yield very good results in an itraconazole formulation both as regards physico-chemical stability as well as regards organoleptic acceptability. In the high-dosage formulations 35 stronger flavours are required such as Caramel Chocolate flavour, Mint Cool flavour, Fantasy flavour and the like pharmaceutically acceptable strong flavours. Each flavour may be present in the final composition in a concentration ranging from 0.05% to 1%

-6-

(w/v). Combinations of said strong flavours are advantageously used. Preferably a flavour is used that does not undergo any change or loss of taste and colour under the acidic conditions of the formulation.

- 5 A preferred high-dosage formulation according to the present invention comprises by weight or by volume based on the total volume of the formulation :
- (a) 4% (w/v) itraconazole;
 - (b) 60% (w/v) hydroxypropyl- β -cyclodextrin;
 - (c) 10% (v/v) propylene glycol;
 - 10 (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
 - (e) 0.08% (w/v) sodium saccharin;
 - (f) up to 1% (w/v) of one or more strong flavours; and
 - (g) water.
- 15 The preparation of the formulations according to the present invention will hereafter be described with regard to a preferred low-dosage formulation having the following composition (% are by weight or by volume based on the total volume of the formulation) :
- (a) 1% (w/v) itraconazole;
 - 20 (b) 40% (w/v) hydroxypropyl- β -cyclodextrin;
 - (c) 10% (v/v) propylene glycol;
 - (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
 - (e) 0.06% (w/v) sodium saccharin;
 - (f) 19% (v/v) sorbitol (70%) non-crystallizing solution;
 - 25 (g) up to 1% (w/v) of one or more cherry flavours; and
 - (h) water.

Optionally, the above preferred low-dosage formulation further comprises up to 0.1%, in particular 0.02% caramel sweetener.

30

Similar formulations can be prepared with saperconazole, though other flavours may be preferred then.

Said process of preparation comprises the steps of

- 35 (a) dissolving the active ingredient in the alcoholic co-solvent and acid;
- (b) dissolving the cyclodextrin in water and adding thereto the solution prepared in (a) while stirring until homogenous;

-7-

- (c) adding the sweetener(s) and the flavour(s);
- (d) adjusting the acidity to $\text{pH } 2.0 \pm 0.1$ and
- (e) diluting the formulation to the desired end-volume.

- 5 In particular, for preparing 1 litre of the aforementioned preferred formulation 100 ml of propylene glycol is treated with 3.76 ml concentrated HCl, stirred and slightly heated. 10 g itraconazole is added and stirring is continued until homogeneous. In a separate vessel, 400 g hydroxypropyl- β -cyclodextrin is dissolved in 400 ml distilled water. The solution of the active ingredient is added slowly to the cyclodextrin solution
- 10 while stirring. The sorbitol solution (190 ml) is added and stirred till homogeneous. The sodium saccharin (0.6 g) is dissolved in 50 ml distilled water and added to the mixture. The flavours are added and the pH of the mixture (about 1.7) is adjusted with a 10 N NaOH solution to $\text{pH } 2.0 \pm 0.1$. The resulting solution is diluted with distilled water to an end volume of 1 litre. A pharmaceutical dosage form is obtained by filtering
- 15 the previous solution and filling it into suitable containers. e.g. in 100 ml glass bottles with a screw cap. The pharmaceutical dosage form advantageously comprises a minimal volume of air above the solution, preferably an inert gas such as nitrogen. Besides the exclusion of air (oxygen), storage at temperatures below 25°C also beneficially affects the maximum shelf life of the formulation for oral administration.
- 20 In case a more simple formulation lacking the flavour(s) and/or sweetener(s) is envisaged, step (c) is omitted partially or completely from the process of preparation.

Claims

- 5 1. A formulation for oral administration comprising an antifungal, a sufficient amount of a cyclodextrin or a derivative thereof, an aqueous acidic medium as bulk liquid carrier and an alcoholic co-solvent.
- 10 2. A formulation according to claim 1 further comprising one or more pharmaceutically acceptable sweeteners and one or more pharmaceutically acceptable flavours.
- 15 3. A formulation according to claim 1 or 2 wherein the antifungal is itraconazole or saperconazole and the cyclodextrin is hydroxypropyl- β -cyclodextrin having an M.S. in the range of 0.35 to 0.50 and containing less than 1.5% unsubstituted β -cyclodextrin.
- 20 4. A formulation according to claim 3 wherein the alcoholic co-solvent is propylene glycol.
5. A formulation according to claim 4 having a pH of 2.0 ± 0.1 .
6. A formulation according to claim 5 wherein the pharmaceutically acceptable sweetener comprises at least one intense sweetener and optionally a bulk sweetener.
- 25 7. A formulation according to claim 6 wherein the intense sweetener is selected from the group consisting of saccharin, sodium or calcium saccharin and the bulk sweetener is selected from the group consisting of sorbitol, mannitol, fructose, sucrose, maltose, glucose, caramel or honey.
- 30 8. A formulation according to claim 2 comprising by weight or by volume based on the total volume of the formulation :
- 35 (a) 4% (w/v) itraconazole;
- (b) 60% (w/v) hydroxypropyl- β -cyclodextrin;
- (c) 10% (v/v) propylene glycol;
- (d) acid and base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
- (e) 0.08% (w/v) sodium saccharin;
- (f) up to 1% (w/v) of one or more flavours; and
- (g) water.

9. A formulation according to claim 2 comprising by weight or by volume based on the total volume of the formulation :
- (a) 1% (w/v) itraconazole or saperconazole;
 - 5 (b) 40% (w/v) hydroxypropyl- β -cyclodextrin;
 - (c) 10% (v/v) propyleneglycol;
 - (d) acid or base to adjust the pH of the composition within the range of 2.0 ± 0.1 ;
 - (e) 0.06% (w/v) sodium saccharin;
 - (f) 19% (v/v) sorbitol (70%) non-crystallizing solution;
 - 10 (g) up to 1% (w/v) of one or more flavours;
 - (h) 0.02% (w/v) of a caramel sweetener; and
 - (i) water.
10. A process of preparing a formulation as claimed in claim 1, characterized in that
- 15 said process comprises the steps of :
- (a) dissolving the active ingredient in the alcoholic co-solvent and acid;
 - (b) dissolving the cyclodextrin in water and adding thereto the solution prepared in
 - (a) while stirring until homogenous;
 - (c) adding the sweetener(s) and the flavour(s), if any;
 - 20 (d) adjusting the acidity to pH 2.0 ± 0.1 and
 - (e) diluting the formulation to the desired end-volume.

INTERNATIONAL SEARCH REPORT

 Internat Application No
 PCT/EP 94/03169

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K31/495 A61K9/08 A61K47/40 A61K47/10

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANTIMICROB. AGENTS CHEMOTHER., vol.36, no.2, February 1992 pages 477 - 480 J.S. HOSTETLER ET AL. 'Effect of cyclodextrin on the pharmacology of antifungal oral azoles' * see especially p. 478 right column - p. 479 left column *	1,3-5,10
Y	* " *	2,6
P,X	--- WO,A,93 19061 (JANSSEN) 30 September 1993 * see claims 1-3,5-12, p. 11 line 26 - p. 13 line 25 *	1
Y	--- US,A,4 916 134 (HEERES ET AL.) 10 April 1990 cited in the application * see especially examples 12 and 13 *	2,6
	--- -/--	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 January 1995

Date of mailing of the international search report

1 0.02 95

Name and mailing address of the ISA

 European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
 Fax: (+31-70) 340-3016

Authorized officer

Isert, B

INTERNATIONAL SEARCH REPORT

Internat . Application No
PCT/EP 94/03169

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>INT. J. PHARMACEUT., vol.80, 1992 pages 253 - 258 J. PITHA ET AL. 'Preparation of drug: hydroxypropylcyclodextrin complexes by a method using ethanol or aqueous ammonium hydroxide as co-solubilizer' * see especially summary and p. 258 *</p> <p>-----</p>	1

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat Application No

PCT/EP 94/03169

Patent document cited in search report	Publication date	Patent family member(s)		Publication date
WO-A-9319061	30-09-93	AU-B-	3632493	21-10-93
		CA-A-	2117651	30-09-93
		EP-A-	0631578	04-01-95
		FI-A-	944311	16-09-94
		NO-A-	943450	11-11-94

US-A-4916134	10-04-90	AU-B-	600107	02-08-90
		AU-A-	1358588	29-09-88
		DE-A-	3874576	22-10-92
		DK-B-	168336	14-03-94
		EP-A, B	0283992	28-09-88
		ES-T-	2044991	16-01-94
		JP-B-	6067929	31-08-94
		JP-A-	63277674	15-11-88
		SU-A-	1635900	15-03-91
